GENERAL INDEX TO VOLUME XXXVI

New scientific names of plants and the final members of new combinations are printed in **bold-face** type; synonyms and page numbers having reference to figures and plates, in *italics;* and all other matter in ordinary type.

A

Adenodiscus, 511; *mexicanus,* 523
Allen, P. H. Orchidaceae of Panama (third part), 1; (fourth part), 133
Ancash, On some uses of maize in the Sierra of, 405
Anderson, Edgar: Charles M. Rick and. On some uses of maize in the Sierra of Ancash, 405; Stoner, C. R., and. Maize among the Hill Peoples of Assam, 355
Andrews, Henry N., Jr. Nucellangium, a new genus of fossil seeds previously assigned to Lepidocarpon, 479
Aneurophyton, 291
Apocynaceae of South America, New, 543
Asia: maize in, 389; distribution of major races in, 391
Assam, Maize among the Hill Peoples of, 355
Assam hills: economics in the, 375; history in the, 374; map of, *357;* tribes of, 356; varieties of maize among the, 359
Azumbuja, David de, and Robert E. Woodson, Jr. New Apocynaceae of South America, 543

B

Baxter, Robert W. Some Pteridosperm stems and fructifications with particular reference to the Medullosae, 287
Beer prepared from maize: in Assam, 377, in Peru, 410
Bhutan, maize in, 372
Burma, maize in, 373

C

Calamopitys, 291
California, Assamese varieties of maize grown in, 356, 379
Callejon de Huailas, uses of maize in the, 406
Cancha, preparation of, from maize, 409
Carboniferous plants, 287, 479
Cardiocarpus, 492; bicuspidatus, 493; injens, 492; minor, 492
Chicha, preparation of, from maize, 410; chicha de jora, 410; chicha morada, 411
Chochaca, preparation of, from maize, 411
Chondrorhyncha **aromatica,** 85; **caloglossa,** 85; **costaricensis,** 86; **discolor,** 87; **marginata,** 88, *89*
Chromosomes in Paphiopedilum: comparative length of, in P. Lawrenceanum and P. callosum, *441, 443;* numbers of, 445; terminal, 453, *468*
Coal balls, fossils found in, 287, 479
Coix, uses of, by the Assamese tribes, 358
Cordaianthus spicatus, 493
Cusqueño blanco, 406
Cypripedilum, 433
Cypripedium, 433; *callosum,* 434; *Hyeanum,* 435; *Lawrenceanum* var. *Hyeanum,* 435
Cytology: of Paphiopedilum callosum, 447; of P. Maudiae Hort., 433, 449; of P. Lawrenceanum, 448; analysis of root-tips, 440; of yeasts, 260

D

Dafla tribe of Assam, maize culture by the, 368, *398*
Darrah, W. H., description of coal-ball fossil by, 479
Dolerotheca, 315; formosa, 316, *344, 348;* Reedana, 317; **Schopfii,** 317, *348, 350, 352;* **sclerotica,** 315, *316, 346, 348, 352;* villosa, 317

E

Economics in the Assam hills, 375
Euspermopteris, 294
Ethnology in relation to maize, 355, 405

F

Ferns, fossil, 287
Food plants among the Hill Peoples of Assam, 374
Fossil: fructifications, 315; seeds, 479; stems, 287

G

Galactose, fermentation of, 260, 265
Grewia terebinthinaceus, 513

H

Heliocarpus, A revision of the genus, 507; generic relationships, 507; economic value of, 509; geographic distribution, 509; taxonomy, 511; vernacular names, 510
Heliocarpus, 511; americanus, 528, 532; *americanus* var. *popayanensis*, 532, var. *typica*, 528; appendiculatus, 526; attenuatus, 520; *australis*, 532; *belizensis*, 523; *boliviensis*, 532; *costaricensis*, 523; *Caecaliae*, 530; *chontalensis*, 526; *cuspidatus*, 530; *diclinus*, 532; Donnell-Smithii, 530; *excelsior*, 536; *floribundus*, 530; *Gentlei*, 536; *glaber*, 517; *glabrescens*, 523; *glanduliferus*, 523; *horridus*, 530; *laevis*, 521; mexicanus, 523; *microcarpus*, 513; *Nelsoni*, 513; nodiflorus, 536; occidentalis, 521; pallidus, 515; Palmeri, 517; *polyandrus*, 517, var. *nodiflorus*, 536; popayanensis, 532; *popayanensis* var. *grandifolius*, 532, var. *Schumanni*, 532, var. *trichopoda*, 532; *reticulatus*, 513; *Rosei*, 532; *rudis*, 532; *stipulatus*, 532; *subtrilobus*, 532; terebinthinaceus, 513; *tomentosus*, 528; *trichopodus*, 532; *velutinus*, 515; *viridis*, 520; *yucatanensis*, 523
Hermann, Frederick J. New species of Lonchocarpus from Panama, 281
Heterangium Grievii, 204; Kukuki, 294; minimum, 294
Hybrids: of Paphiopedilum, 433; of Saccharomyces, 259

I

Inheritance in yeast, 260
Internode diagram of a variety of Assamese maize, *381*
Iowa, fossils found in coal-balls from, 287, 479

J

Job's tears, use of, by the Assamese tribes, 358

K

Kaloxylon, 293

L

Lacmellea **speciosa**, 543
Lay, Ko Ko. A revision of the genus Heliocarpus L., 507
Lepidocarpon, 486; glabrum, 479, 491; lomaxi, 486
Lonchocarpus, New species of, from Panama, 281
Lonchocarpus atropurpureus, 282; **calcaratus**, 282; densiflorus, 284; glabrescens, 284; **oliganthus**, 281; parviflorus, 282; praecox, 284
Lyginopteris, 290

M

McQuade, Henry A. Cytology of Paphiopedilium Maudiae Hort., 433
Maíz dulce, 409; maíz morada, 411
Maize: among the Hill Peoples of Assam, 355, culture of, 376; geographic distribution of, 405; history of, 405; in hill areas outside Assam, 372; in the Orient, 355; morphological survey of, grown by the Nagas, 379; On some uses of, in the Sierra of Ancash, 405; races of, in South America and Asia, 390; uses of, in Assam, 377, in Peru, 405; varieties of, 380, 383
Maize: "Caribbean", 383; "Drooping Waxy", 381, 386, *402*; "Early Slender", 383, *400*; "Early Upright", 384, *404*; "Late Sidewise", *381*, 385, *402*; "Late Upright", 383, 384, *404*.
Malouetia cestroides, 544; **grandiflora**, 544; **parviflora**, 544
Manioc, culture of, in the Assam Hills, 374
Manipur State: maize in, 373; tribes of, 357
Marsilea, 492
Medicine, use of corn preparations in, 409
Medullosa, 290; anglica, var. ioensis, 298; centrofilis, 298; distelica, 298; **elongata**, 301; **endocentrica**, 303; Leuckarti, 298; Noei, 298; **primaeva**, 298; Thompsonii, 298
Medullosae, Some Pteridosperm stems and fructifications with particular reference to the, 287
Melibiose, fermentation of, 260, 265
Mendelian, Non-, segregations in Saccharomyces, 259
Microspermopteris aphyllum, 289, *322*, *324*; partial reconstruction of, *382*
Millet, use of, by the Assamese tribes, 358
Monba tribe of Assam, 358, *396*; threshing maize, *396*; maize stacked for ripening, *396*
Mote, preparation of, from maize, 411
Mundkur, Balaji D. Evidence excluding mutations, polysomy, and polyploidy as possible causes of non-Mendelian inheritance in Saccharomyces, 259

N

Naga tribes of Assam, 357, 359, *398*; corner of a "jhum" field, *398*; varieties of maize

grown by, 379, characteristics of, 384
Nepal, maize in, 373
Non-Mendelian segregations in Saccharomyces, Evidence as possible causes of, 259
Nucellus, fossil, 480, 486
Nucellangium, a new genus of fossil seeds previously assigned to Lepidocarpon, 479
Nucellangium glabrum, 491

O

Olla canchera, a prehistoric vessel used to parch maize, 409, *409*
Orient, maize in the, 355
Orchidaceae: of Panama (third part), 1, (fourth part), 133; Index (part III), 247; Pfitzer's key to Pleonandrae of, 434; Rolfe's key to suborder Diandrae of, 434

P

Paleobotany studies, 287
Panama: Flora of, Part III, Fascicle 4 (Orchidaceae, third part), 1, Fascicle 5 (Orchidaceae, fourth part), 133, Index to Part III, 247; New species of Lonchocarpus from, 281; A first record of the genus Qualea in, 285
Paphiopedilum Maudiae Hort., The cytology of, 433
Paphiopedilum, 433; albino forms of, 437; chromosome numbers in, 445; cytology of P. callosum, 447, of P. Lawrenceanum, 448, of P. Maudiae, 449; key to subgenera, 435; meiosis in, 447; root-tips, 440; taxonomy of, 433
Paphiopedilum: "Alma Gavaert", 438; bellatulum, var. album, 437; callosum, 434, 437, 441, 447, *466*, list of crosses in which used as parent, 459, var. Sanderae, 433, 434, 437, 459; Charlesworthii var. album, 437; Curtisii, 437; "Emerald", 438; "Enchantress", 438; Holdenii, 438; insigne var. Sanderae, 438, var. Sanderianum, 438; Lawrenceanum, 435, 437, 448, *466*, list of crosses in which used as parent, 462, var. Hyeanum, 433, 435, 437; Maudiae, 433, 435, analysis of, 448, list of crosses in which used as parent, 456, var. "Bank House", 436, var. "Dell", 436, var. magnificum, 436, *466*, var. "Westonbirt", 436; niveum, 437; Rosettii, 438; "Warden", 438
Paphiopedium, 433
Peru, uses of maize in Ancash, Dept. of, 405
Psilophyton princeps, 295
Photographing spores of Selaginella, 414
Phragmopedilum, 433
Phragmopedium, 433
Phytolaccaceae, 475
Pollen grains: of Paphiopedilum callosum, 447, *470*; of P. Lawrenceanum, 449, *470*; of P. Maudiae, 450, *470*
Polynesians, transfer of crops by, 292
Popcorns: in Assam, 379; Oriental, 355; in Peru, 409
Potato, culture of, in the Assam Hills, 374
Pteridosperm stems and fructifications, Some, with particular reference to the Medullosae, 287

Q

Qualea, A first record for the genus (Vochysiaceae), in North America, 285
Qualea **cymulosa**, 285; rupicola, 286

R

Rhetinangium, 290
Rice in the Assam Hills, 358
Rick, Charles M., and Edgar Anderson. On some uses of maize in the Sierra of Ancash, 405
Rogers, David J. Stegnosperma: A new species and a generic commentary, 475

S

Saccharomyces: Evidence excluding mutations, polysomy and polyploidy as possible causes of non-Mendelian inheritance in, 259; tetrad analyses of crosses of, 260
Schery, Robert W.: A first record for the genus Qualea (Vochysiaceae) from North America (Panama), 285; Robert E. Woodson, Jr., and collaborators. Flora of Panama, Part III, Fasc. 4, 1, Fasc. 5, 133
Schopfiastrum, 291
Secondatia **Adolphii**, 543, *546*, *548*; Duckei, 543
Seed-ferns, 288; comparative characters of stem genera of, 290
Seeds, fossil, 479, "normal", 480, "proliferated", 486
Segregations in yeast, Non-Mendelian, 259, 271
Selaginella, Spores of the genus, in North America north of Mexico, 413
Selaginella, 413; acanthonota, 419, *426*; apoda, 423, *431*; arenicola, 418, *425*; arizonica, 422, *430*; armata, 413, 414, 422, *431*; asprella, 420, 421, *429*; Bigelovii, 417, 418, *425*; cinerascens, 419, *427*; Ser. Circinatae, 423; Coryi, 418; Ser. Decumbentes, 422; densa, 413, 420, 421, *429*; Douglasii, 422, *431*; eremophila, 422, *430*; subg. Euselaginella, 415, 417;

floridana, 419, *426*; funiformis, 419, *426*; Hanseni, 421, 422, *430*; lepidophylla, 423, *431*; leucobryoides, 420, *428*; mutica, 419, *427*; neomexicana, 418; oregana, 419, *426*; Parishii, 422; pilifera, 413, 423, *431*; Riddellii, 418, *425*; rupestris, 413, 417, 420, 421, *429*; rupincola, 417, 418, *425*; scopulorum, 420, 421, *429*; selaginoides, 413, 414, 415, 417, *424*; Sheldonii, 422, *430*; *Sherwoodii*, 417, *424*; sibirica, 421, *428*; subg. Stachygynandrum, 416, 422; Standleyi, 420, *428*; tortipila, 414, 417, *424*; Underwoodii, 419, *427*; Wallacei, 413, 421, *428*; Watsoni, 420, *427*; Wrightii, 421, *430*

Selenipedilum, 433

Selenipedium, 433

Sierra of Ancash, uses of maize in, 406

Sikkim, maize in, 372

Sorghum: relation between maize and, 393; use of, by the Assamese tribes, 358

South America, New Apocynaceae of, 543; maize of, 389, 405

Sporangium, fossil, 480, 486

Spores of the genus Selaginella in North America north of Mexico, 413

"Star rings" from carboniferous species of Medullosa, 312, *338*, *340*, *342*

Stegnosperma: A new species and a generic commentary, 475; cubense, 476; *halimifolia*, 477; halimifolium, 475, 477; *halimifolium*, 476; *scandens*, 476; Watsonii, 475

Stem genera of seed-ferns, 290

Stems and fructifications, Some Pteridosperm, with particular reference to the Medullosae, 287

Stenomylon, 291

Stonor, C. R., and Edgar Anderson. Maize among the Hill Peoples of Assam, 355

Sutcliffia, 291

T

Tercio pelo, 409

Tetrad analyses of yeast hybrids, 260, 268

Tiestos, 409

Tocos, preparation of, from maize, 406

Trichilia scandens, 476

Triumfetta, 1; *mexicana*, 523

Tryon, Alice F. Spores of the genus Selaginella in North America north of Mexico, 413

V

Valcárcel's illustration of Olla canchera, 409

Vitamin synthesizing abilities in yeasts, segregations of, 271

Vochysiaceae, the genus Lonchocarpus of the, 285

W

Woodson, Robert E., Jr., and David de Azumbuja. New Apocynaceae of South America, 543; Robert W. Schery and, and collaborators. Flora of Panama, Part III, Fasc. 4, 1, Fasc. 5, 133

Y

Yeast hybrids, tetrad analysis of, 260

Yeast, see Saccharomyces

Z

Zea Mays—see Maize